

A Gravity Theory on Noncommutative Spaces

Frank Meyer

Max-Planck Institute for Physics Munich

Based on joined work with:

P. Aschieri, C. Blohmann, M. Dimitrijević, P. Schupp and J. Wess

see preprint **hep-th/0504183**



Outline

- Deformed Spaces and Symmetries
- Bialgebra of Deformed Diffeomorphisms
- Representations: Scalar, Vector and Tensor Fields
- Metric and Christoffel-Symbol
- Covariant Derivatives and Curvature
- θ -Deformed Einstein-Hilbert action
- Outlook



Deformed Spaces

- **Underlying idea:** Discrete Spacetime
At very short distances: coordinates do not commute
(Heisenberg 1930)

$$[\hat{x}^\mu, \hat{x}^\nu] = i\theta^{\mu\nu}(\hat{x}) \neq 0$$

- differentiable space-time manifold \longrightarrow algebra of noncommutative coordinates:

$$\hat{\mathcal{A}}_{\theta(x)} := \mathbb{C}\langle\langle \hat{x}^1, \dots, \hat{x}^n \rangle\rangle / ([\hat{x}^\mu, \hat{x}^\nu] - \theta^{\mu\nu}(\hat{x}))$$

- We consider elements of $\hat{\mathcal{A}}_{\theta(x)}$ as “functions” on the deformed Space
- Easiest example: θ -Deformed or Canonically Deformed Quantum Space $\hat{\mathcal{A}}_\theta$

$$[\hat{x}^\mu, \hat{x}^\nu] = i\theta^{\mu\nu}, \quad \theta^{\mu\nu} \in \mathbb{R}$$



Star Product Realization

- Vector spaces of formal power series in commutative coordinates and noncommutative coordinates are isomorphic (*Poincaré-Birkhoff-Witt property*)

$$\begin{aligned}\rho : \mathbb{C}[x^0, \dots, x^n][[\theta]] &\rightarrow \hat{\mathcal{A}}_\theta \\ f(x^\mu) &\mapsto \hat{f}(\hat{x}^\mu)\end{aligned}$$

- We define a *new product*, called **star product** by pulling back the product of the noncommutative algebra:

$$f(x^\mu) \star g(x^\mu) := \rho^{-1}(\hat{f}(\hat{x}^\mu) \cdot \hat{g}(\hat{x}^\mu))$$

- This way we realized the deformed algebra $\hat{\mathcal{A}}_\theta$ on the algebra of commutative functions equipped with a new (*noncommutative*) product \star



Star Product Realization

- The star product is *not* unique since the isomorphism ρ (*ordering prescription*) is not unique
- *Example:* Canonical Structure $[\hat{x}^\mu, \hat{x}^\nu] = i\theta^{\mu\nu}$
Moyal-Weyl Product

$$f \star g = \mu \circ e^{\frac{i}{2}\theta^{\mu\nu}\partial_\mu \otimes \partial_\nu} (f \otimes g) = fg + \frac{i}{2}\theta^{\mu\nu}(\partial_\mu f)(\partial_\nu g) + \dots,$$

where $\mu(f \otimes g) := fg$ is just the multiplication map and

$$[x^\mu \star, x^\nu] = i\theta^{\mu\nu}$$

- For $\theta \rightarrow 0$ we get back the commutative product
- The star product above is associative



Derivatives

- **Derivatives** are maps on the deformed coordinate space:

$$\hat{\partial}_\mu : \hat{\mathcal{A}}_\theta \rightarrow \hat{\mathcal{A}}_\theta$$

- Thus, they have to be consistent with the commutation relations of the coordinates.
- general ansatz:

$$[\hat{\partial}_\mu, \hat{x}^\nu] = \delta_\mu^\nu + \sum_j A_\mu^{\nu\rho_1 \dots \rho_j} \hat{\partial}_{\rho_1} \dots \hat{\partial}_{\rho_j}.$$

- θ -Deformed Space: The *undeformed* calculus

$$[\hat{\partial}_\mu, \hat{x}^\nu] := \delta_\mu^\nu$$

is consistent with the algebra relations $[\hat{x}^\mu, \hat{x}^\nu] = i\theta^{\mu\nu}$



★-Action and θ -Deformed Diffeomorphisms

- **Diffeomorphisms** are generated by vector fields $\xi = \xi^\mu \partial_\mu$ which satisfy the Lie algebra

$$[\xi, \eta] = \xi \times \eta = (\xi^\mu (\partial_\mu \eta^\nu) - \eta^\mu (\partial_\mu \xi^\nu)) \partial_\nu$$

- Aim: Lift vector fields to differential operators which define maps on the algebra \hat{A}_θ , i.e. are consistent with the commutation relations of the coordinates
- Since we have a well-defined differential calculus also the following map

$$\hat{\xi} : \hat{f} \mapsto \hat{\xi}^\rho \hat{\partial}_\rho \hat{f}$$

is a well-defined map on the deformed space \hat{A}_θ

- Mapped to the functions of commuting variables we obtain the **★-action**:

$$\xi : f \mapsto \xi^\rho \star (\partial_\rho f) =: \xi \triangleright f$$



★-Action and θ -Deformed Diffeomorphisms

But: The \star -commutator of two such operators does not close and we don't have a representation of the Lie algebra of vector fields:

$$[\xi \star \eta] = \xi \times \eta = (\xi^\mu \star (\partial_\mu \eta^\nu) - \eta^\mu \star (\partial_\mu \xi^\nu)) \partial_\nu + [\xi^\mu \star \eta^\nu] \partial_\mu \partial_\nu$$

- Define higher order differential operator by requiring

$$[X_\xi \star X_\eta] = X_{[\xi, \eta]}$$

where $\xi = \xi^\rho \partial_\rho$ is an ordinary vector field. Then the operators X_ξ have the Lie algebra structure of vector fields!

- Solution: $X_\xi = \xi - \frac{i}{2} \theta^{\mu\nu} (\partial_\mu \xi) \partial_\nu + \dots$
- Mapped to the space of commutative coordinates we get

$$X_\xi \triangleright f = \xi f = \xi^\mu (\partial_\mu f)$$



★-Action and θ -Deformed Diffeomorphisms

- The operators X_ξ act in a nontrivial way on the \star -product of two functions:

$$X_\xi \triangleright (f \star g) = (X_\xi \triangleright f) \star g + f \star (X_\xi \triangleright g) - \frac{i}{2} \theta^{\rho\sigma} \{((\partial_\rho X_\xi) \triangleright f) \star (\partial_\sigma g) + (\partial_\rho f) \star ((\partial_\sigma X_\xi) \triangleright g)\} + O(\theta^2)$$

- One can abstract this deformed Leibniz rule to a nontrivial **coproduct**

$$\Delta X_\xi = X_\xi \otimes 1 + 1 \otimes X_\xi - \frac{i}{2} \theta^{\rho\sigma} \{((\partial_\rho X_\xi) \otimes \partial_\sigma + \partial_\rho \otimes (\partial_\sigma X_\xi))\} + O(\theta^2)$$

- The coproduct tells us how to act on a product of two functions



Scalar, Vector and Tensor Fields

- We introduce scalar fields to be elements ϕ in $\hat{\mathcal{A}}_\theta$ with the following transformation property:

$$\delta_\xi \phi = -X_\xi \triangleright \phi$$

- The coproduct

$$\Delta \delta_\xi = \delta_\xi \otimes 1 + 1 \otimes \delta_\xi - \frac{i}{2} \theta^{\rho\sigma} \{((\partial_\rho \delta_\xi) \otimes \partial_\sigma + \partial_\rho \otimes (\partial_\sigma \delta_\xi))\} + O(\theta^2)$$

assures that a \star -product of scalar fields transforms as a scalar again:

$$\delta_\xi(\phi \star \psi) = -X_\xi \triangleright (\phi \star \psi)$$

- Similarly we define vector fields to transform like

$$\delta_\xi V_\mu = -X_\xi \triangleright V_\mu - X_{\xi^\rho} \star V_\rho$$

and tensor fields in full analogy



Summary: θ -Deformed Diffeomorphisms

- To summarize, we end up with the **θ -Deformed Diffeomorphisms:**

The algebra is undeformed

$$[\delta_\xi, \delta_\eta] = \delta_{\xi \times \eta}$$

but the coalgebra is deformed

$$\Delta(\hat{\delta}_\xi) = e^{-\frac{i}{2}\theta^{\rho\sigma}\partial_\rho^* \otimes \partial_\sigma^*} \left(\hat{\delta}_\xi \otimes 1 + 1 \otimes \hat{\delta}_\xi \right) e^{\frac{i}{2}\theta^{\rho\sigma}\partial_\rho^* \otimes \partial_\sigma^*}$$

- We have scalar fields and tensor fields as representations of it
- On scalar fields the above deformed algebra is represented by the differential operators X_ξ from the previous discussion



Metric and Christoffel-Symbol

- We define the metric to be a symmetric tensor of rank two which reduces to the classical metric $g_{\mu\nu}$ for $\theta \rightarrow 0$
- This can be done using a set of four vectors E_μ^a (**Vierbeins**)

$$G_{\mu\nu} = \frac{1}{2} \left(E_\mu^a \star E_\nu^b + E_\nu^a \star E_\mu^b \right) \eta_{ab},$$

where η_{ab} is the usual metric of flat Minkowski space

- The deformed coproduct assures that $G_{\mu\nu}$ transforms indeed as a tensor
- To raise and lower indices we need its \star -inverse $G^{\mu\nu\star}$

$$G_{\mu\nu} \star G^{\nu\rho\star} = \delta_\mu^\rho$$



Metric and Christoffel-Symbol

- This is again a tensor in \mathcal{A}_θ and can be constructed in terms of the classical inverse $G^{\mu\nu}$
- Let us assume that the connection is symmetric
 $\Gamma_{\mu\nu}^\alpha = \Gamma_{\nu\mu}^\alpha$
- Demanding that the covariant derivative of $G_{\mu\nu}$ vanishes

$$D_\alpha G_{\beta\gamma} = \partial_\alpha^\star \triangleright G_{\beta\gamma} - \Gamma_{\alpha\beta}^\rho \star G_{\rho\gamma} - \Gamma_{\alpha\gamma}^\rho \star G_{\beta\rho} = 0$$

we obtain $\Gamma_{\mu\nu}^\alpha$ uniquely expressed in terms of $G_{\mu\nu}$ and $G^{\mu\nu}$ **(Christoffel-Symbol)**

$$\Gamma_{\alpha\beta}^\sigma = \frac{1}{2} \left(\partial_\alpha^\star \triangleright G_{\beta\gamma} + \partial_\beta^\star \triangleright G_{\alpha\gamma} - \partial_\gamma^\star \triangleright G_{\alpha\beta} \right) \star G^{\gamma\sigma}$$



Covariant Derivatives and Curvature

- We are able to construct covariant derivatives

$$\mathcal{D}_\mu V_\nu := \partial_\mu V_\nu - \Gamma_{\mu\nu}^\alpha \star V_\alpha$$

such that $\mathcal{D}_\mu V_\nu$ transforms as a tensor

- The curvature tensor can be obtained as the commutator of two covariant derivatives

$$[\mathcal{D}_\mu \star, \mathcal{D}_\nu] V_\rho = R_{\mu\nu\rho}^\sigma \star V_\sigma + T_{\mu\nu}^\alpha \star \mathcal{D}_\alpha V_\rho$$

- The curvature can be expressed in terms of the connection and can be expanded in orders of θ :

$$\begin{aligned} R_{\mu\nu\rho}^\sigma &= R_{\mu\nu\rho}^{(0)\sigma} - \frac{i}{2} \theta^{\kappa\lambda} (\partial_\kappa R_{\mu\nu\rho}^{(0)\tau}) (\partial_\lambda g_{\tau\gamma}) g^{\gamma\delta} \\ &\quad + \frac{i}{2} \theta^{\kappa\lambda} (\partial_\kappa \Gamma_{\nu\rho}^{(0)\beta}) (\Gamma_{\mu\beta}^{(0)\tau}) (\partial_\lambda g_{\tau\gamma}) g^{\gamma\sigma} - \Gamma_{\mu\tau}^{(0)\sigma} (\partial_\lambda g_{\beta\gamma}) g^{\gamma\tau} + (\partial_\lambda \Gamma_{\mu\beta}^{(0)\sigma}) \\ &\quad - \frac{i}{2} \theta^{\kappa\lambda} (\partial_\kappa \Gamma_{\mu\rho}^{(0)\beta}) (\Gamma_{\nu\beta}^{(0)\tau}) (\partial_\lambda g_{\tau\gamma}) g^{\gamma\sigma} - \Gamma_{\nu\tau}^{(0)\sigma} (\partial_\lambda g_{\beta\gamma}) g^{\gamma\tau} + (\partial_\lambda \Gamma_{\nu\beta}^{(0)\sigma}) + \dots, \end{aligned}$$

where $R_{\mu\nu\rho}^{(0)\sigma}$ is the commutative curvature tensor



Deformed Einstein-Hilbert Action

- We can define the **θ -deformed Einstein-Hilbert action**

$$S_{\text{EH}} = \frac{1}{2} \int d^4x \left(E^* \star R + \text{c.c.} \right)$$

- It is real by definition and invariant with respect to the deformed diffeomorphisms
- Since the integral is cyclic

$$\int d^4x f \star g = \int d^4x g \star f$$

we can write (if the measure E^* is real):

$$S_{\text{EH}} = \frac{1}{2} \int d^4x \left(E^* \star R + \bar{R} \star E^* \right) = \frac{1}{2} \int d^4x E^* \star (R + \bar{R})$$



Discussion and Outlook

- We have constructed a θ -Deformed Bialgebra of Diffeomorphisms.
- We have studied scalar, vector and tensor fields as representations of this θ -Deformed Bialgebra of Diffeomorphisms.
- We introduced a covariant derivative and the curvature tensor
- We defined a metric and got the curvature scalar
- We construct a θ -deformed diffeomorphism invariant Einstein-Hilbert action and got

A Gravity Theory on Deformed Spaces

... see hep-th/0504183 for more details

